



3425 Corporate Way Duluth, GA 30096



Patient: SAMPLE PATIENT DOB: Sex: MRN:

3301 Organix ® Comprehensive Profile - Urine *Methodology: LC/Tandem Mass Spectrometry, Colorimetric*

Summary of Abnormal Findings				
Biomarkers	Findings	Metabolic Pathway		
Fatty Acid Metabolism				
Adipate	Borderline High	Fatty acid oxidation		
Suberate	Borderline High	Fatty acid oxidation		
Carbohydrate Metabolism				
L-Lactate	Н	Glycolysis		
b-Hydroxybutyrate	Borderline High	Ketone production		
Energy Production Markers				
Succinate	Borderline High	Citric acid cycle		
B-Complex Vitamin Markers	No Abnormality Found			
Methylation Cofactor Markers	No Abnormality Found			
Neurotransmitter Metabolism Markers				
Vanilmandelate	Borderline High	Epinephrine & norepinephrine metabolism		
Homovanillate	Н	Dopamine metabolism		
5-Hydroxyindoleacetate	Н	Serotonin metabolism		
Kynurenate	Borderline High	Tryptophan pathway		
Oxidative Damage and Antioxidant Markers				
p-Hydroxyphenyllactate	Н	Gut bacterial metabolism		
8-Hydroxy-2-deoxyguanosine	Н	Oxidative damage		
Detoxification Indicators				
Sulfate	L	Transsulfuration pathway		
Bacterial - General				

© Genova Diagnostics · Robert M. David, PhD, Lab Director · CLIA Lic. #11D0255349 · Medicare Lic. #34-8475 · Georgia Lab Lic. Code #067-007 New York Clinical Lab PFI #4578 · Florida Clinical Lab Lic. #800008124 Patient: Sample Patient

Patient: Sample Patient	Page 2			
Summary of Abnormal Findings				
Biomarkers	Findings	Metabolic Pathway		
Hippurate	Borderline High	Gut bacterial metabolism		
Phenylacetate	Н	Gut bacterial metabolism		
p-Hydroxybenzoate	Borderline High	Gut bacterial metabolism		
p-Hydroxyphenylacetate	Н	Gut bacterial metabolism		
L. acidophilus/General Bacteria	No Abnormality Found			
Clostridial Species	No Abnormality Found			
Yeast/Fungal				
D-Arabinitol	Borderline High	Yeast product		



3425 Corporate Way Duluth, GA 30096



Patient: SAMPLE PATIENT

DOB: Sex: MRN[.]

3301 Organix® Comprehens	sive Profile - Urine		
Methodology: LC/Tandem Mass Spect This report is not intended for the diagr		of metabolism	
Ranges: Ages 13 and over	Results	QUINTILE DISTRIBUTION	95% Reference
	mcg/mg creatinine	1st 2nd 3rd 4th 5th	Range
		Nutrient Markers	
Fatty Acid Metabolism (Carnitine & B2)		6.2	
1. Adipate	8.5	F F F F F F	┥ <= 11.1
2. Suberate	2.9	2.1	┥ <= 4.6
3. Ethylmalonate	2.0	3.6	┥ <= 6.3
Carbohydrate Metabolism (B1, B3, Cr, Lipoic Acid, CoQ10)	2.0		~= 0.5
4. Pyruvate	<dl< td=""><td>3.9</td><td>┥ <= 6.4</td></dl<>	3.9	┥ <= 6.4
5. L-Lactate	19.0 H	8.5	• 0.6 - 16.4
 β-Hydroxybutyrate 	4.5	2.1 II I I I I ●	┥ <= 9.9
Energy Production (Citric Ac (B Comp., CoQ10, Amino Acids, Mg)	id Cycle)		
7. Citrate	487	601 	56 - 987
8. Cis-Aconitate	38	51	⊣ 18 - 78
9. Isocitrate	62	98	− 39 - 143
	-	19.0	
10. α-Ketoglutarate	<dl< td=""><td>I−−−I−−−−−−−−−−−−−−−−−−−−−−−−−−−−−−</td><td>- <= 35.0</td></dl<>	I −−− I −−−−−−−−−−−−−−−−−−−−−−−−−−−−−−	- <= 35.0
11. Succinate	12.9	I I I I I ↓ ↓ ↓ 0 .59	- <= 20.9
12. Fumarate	<dl< td=""><td></td><td>− <= 1.35</td></dl<>		− <= 1.35
13. Malate	0.6	1.4 	┥ <= 3.1
		3.6	

Page 4

3301 Organix® Comprehensive Profile - Urine

Methodology: LC/Tandem Mass Spectrometry, Colorimetric

This report is not intended for the diagnosis of neonatal inborn errors of metabolism.

Ranges: Ages 13 and over

Ranges: Ages 13 and over		QUINTILE DISTRIBUTION
	Results mcg/mg creatinine	1st 2nd 3rd 4th 5th 95% Reference Range
		Nutrient Markers
B-Complex Vitamin Markers		
(B1, B2, B3, B5, B6, Biotin)		0.25
15. α-Ketoisovalerate	<dl< td=""><td>l = 0.49</td></dl<>	l = 0.49
16. α-Ketoisocaproate	<dl< td=""><td>● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ●</td></dl<>	● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ● ●
	<dl< td=""><td>0.38</td></dl<>	0.38
17. α -Keto- β -Methylvalerate	<dl< td=""><td>ll −−−−−−−−−−−−−−−−−−−−−−−−−−−−−−−−−−−</td></dl<>	l l −−−−−−−−− −−−−−−−−−−−−−−−−−−−−−−−−−−
18. Xanthurenate	<dl< td=""><td>●</td></dl<>	●
		7.6
19. β-Hydroxyisovalerate	3.8	└───┼◆─┼──┼──┤ <= 11.5
Methylation Cofactor Markers (B12, Folate)		
20. Methylmalonate	1.0	1.7
		1.2
21. Formiminoglutamate	0.5	<= 2.2
	Cel	II Regulation Markers
Neurotransmitter Metabolism M		
(Tyrosine, Tryptophan, B6, Antioxidan		1.6 3.9
22. Vanilmandelate	5.2	1 .2 - 5.3
23. Homovanillate	9.1 H	1.9 5.7 1.4 - 7.6
		2.1 5.6
24. 5-Hydroxyindoleacetate	10.5 H	1.6 - 9.8 1.0
25. Kynurenate	1.5	└─── ─────────────────────────────────
		4.0
26. Quinolinate	<dl< td=""><td>◆</td></dl<>	◆
27. Picolinate	3.1	↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓
Oxidative Damage and Antioxid (Vitamin C and Other Antioxidants)	dant Markers	0.39
28. p-Hydroxyphenyllactate	1.13 H	⊢ ⊢ ⊢ ⊢ ⊢ ⊢ ⊢ ⊢ ⊢ ⊢
29. 8-Hydroxy-2-deoxyguanosine	e 13.7 H	5.3 <= 7.6
(Units for 8-hydroxy-2-dexoyguanosine ar	e ng/mg creatinine)	

ID

3301 Organix® Comprehensive Profile - Urine

Methodology: LC/Tandem Mass Spectrometry, Colorimetric

This report is not intended for the diagnosis of neonatal inborn errors of metabolism.

Ranges: Ages 13 and over	Results ncg/mg creatinine	1st	QUI 2nd	NTILE DISTRII 3rd	BUTION 4th	5th	95% Reference Range
	Toxic	ants an	d Detoxi	fication			
Detoxification Indicators (Arg, NAC, Met, Mg, Antioxidants)					0.08	24	
30. 2-Methylhippurate	0.021			-	+ +		<= 0.192
31. Orotate	<dl< td=""><td>•</td><td>1</td><td>-</td><td>0.69</td><td>) </td><td><= 1.01</td></dl<>	•	1	-	0.69) 	<= 1.01
32. Glucarate	<dl< td=""><td>•</td><td>ł</td><td>-</td><td>6.3</td><td></td><td><= 10.7</td></dl<>	•	ł	-	6.3		<= 10.7
33. α-Hydroxybutyrate	<dl< td=""><td>H</td><td></td><td></td><td>0.3</td><td></td><td><= 0.9</td></dl<>	H			0.3		<= 0.9
34. Pyroglutamate	46			+ +	59		28 - 88
35. Sulfate	400 L	958	ł	-		2,347	690 - 2,988
С	compounds o	f Bacter	ial or Ye	ast/Fund	al Origin		
Bacterial - General					0.6		
36. Benzoate	<dl< td=""><td>H</td><td></td><td></td><td></td><td></td><td><= 9.3</td></dl<>	H					<= 9.3
37. Hippurate	570			-	1 1	← 1	<= 1,070
38. Phenylacetate	0.38 H		ł	-	0.1 ²	◆I	<= 0.18
39. Phenylpropionate	<dl< td=""><td>H</td><td></td><td></td><td></td><td></td><td><= 0.06</td></dl<>	H					<= 0.06
40. p-Hydroxybenzoate	1.7				1.1		<= 1.8
41. p-Hydroxyphenylacetate	45 H				19 	I	<= 34
			ľ		64		- 00
42. Indican	32			1	0.73	3	<= 90
43. Tricarballylate	<dl< td=""><td>•</td><td></td><td>-</td><td>1 1</td><td></td><td><= 1.41</td></dl<>	•		-	1 1		<= 1.41
L. acidophilus / General Bacterial					2.0		
44. D-Lactate	0.2	ŀ	- ♦	-	1 1		<= 4.1
Clostridial Species							
45. 3,4-Dihydroxyphenylpropionate	<dl< td=""><td>H</td><td></td><td></td><td></td><td></td><td><= 0.05</td></dl<>	H					<= 0.05
Yeast / Fungal					36		
46. D-Arabinitol	43	 		-	1 1	♦	<= 73
Creatinine = 23 mg/dL							

<DL = less than detection limit

>UL = greater than upper linearity limit

This test has been developed and its performance characteristics determined by Genova Diagnostics, Inc. It has not been cleared by the U.S. Food and Drug Administration.

3301 Organix® Comprehensive Profile - Urine

Additional Considerations

These supplement ranges are not adjusted for age or gender.

Nutrient supplementation is at the *discretion of the treating clinician*. The supplement dose ranges provided below are meant for educational purposes only. These dosage ranges relate to findings commonly found on Genova's nutritional panels and do not apply to specific disease conditions where different dosages may be warranted. Final recommendations should be based on consideration of the patient's medical history and current clinical condition.

Nutrient	Nutrient Need	Clinician Recommendations
Vitamin C	High: 1000-2000 mg	
Vitamin E (mixed tocopherols)	High: 200-400 IU	
Vitamin B-1 (Thiamin)	Low: 10-25 mg	
Vitamin B-2 (Riboflavin)	Low: 10-25 mg	
Vitamin B-3 (Niacin)	Low: 10-50 mg	
Vitamin B-5 (Pantothenic Acid)	Low: 10-25 mg	
Vitamin B-6 (Pyridoxine)	Optional: 0-10 mg	
Magnesium	Optional: 0-100 mg	
Selenium	Optional: 0-50 mcg	
Carnitine	Optional: 0-500 mg	
Coenzyme Q10	Low: 20-60 mg	
Lipoic Acid	Low: 50-100 mg	
N-Acetylcysteine	Low: 100-200 mg	
Need for other antioxidants	High	

Various conditionally essential nurients and other potentially beneficial interventions appear in this section only if relevant abnormalities are present.

Amino acids listed on this page result from functional markers of individual amino acid insufficiency and do not reflect amino acids measured in plasma.

